



Selected Papers of William L. White

www.williamwhitepapers.com

Collected papers, interviews, video presentations, photos, and archival documents on the history of addiction treatment and recovery in America.

Citation: White, W. (2009). Recovery support matching checklist. Posted at www.williamwhitepapers.com

Recovery Support Matching

William L. White

Emeritus Senior Research Consultant
Chestnut Health Systems
bwhite@chestnut.org

Recovery Support Matching

The positive effects of peer-based recovery support groups rise in tandem with dose (the number of meetings attended)(Humphreys, Moos, & Cohen, 1997) and intensity of participation (e.g., applying concepts to daily problem solving, reading recovery literature, sober socializing, service work) (Montgomery, Miller, & Tonigan, 1995). But these positive findings are offset by weak relationships between treatment institutions and local mutual aid groups (White & Hagen, 2005), passive rather than assertive linkage to such groups by addiction professionals (Forman, 2002), and high (approximately 40%) dropout rates from such groups in the first year for adults (Kelly and Moos, 2003) and adolescents (Kelly, Myers, & Brown, 2001). Matching individuals to particular groups and linking individuals to particular persons and meetings can enhance this affiliation process.

The checklist on the following page offers a quick way to assess needs and preferences of particular clients that can help in this matching process. The scales which

follow it provide a way to evaluate a client's intensity of participation in a face-to-face recovery support group program.

Recovery Support Matching Checklist

Which of the following are important to you in selecting a recovery support group? (Check all that apply) I prefer to be with other people who:

- have experience with my primary drug
- are the same gender
- are close to my age
- share my ethnic/cultural background
- share my views on religion, spirituality or secularity
- share my sexual orientation
- smoke tobacco
- do not smoke tobacco
- have tolerant attitudes toward mental illness
- have tolerant attitudes toward medications prescribed for addiction or mental illness
- have prior experience in the criminal justice system

- ___ do not have prior experience in the criminal justice system
- ___ have approximately the same income level
- ___ have had very severe alcohol/drug problems
- ___ have had mild to moderate alcohol/drug problems
- ___ share my goal of complete abstinence
- ___ share my goal of moderated use

Source: Adapted from White, W. & Kurtz, E. (2006). Linking Addiction Treatment and Communities of Recovery: A Primer for Addiction Counselors and Recovery Coaches. Pittsburgh, PA: IRETA/NeATTC

References

- Forman, R. F. (2002). One AA meeting doesn't fit all: 6 keys to prescribing 12-step programs. *Psychiatry Online*, 1(10), 1-6.
- Humphreys, K., Moos, R. J., & Cohen, C. (1997). Social and community resources and long-term recovery from treated and untreated alcoholism. *Journal of Studies on Alcohol*, 58(3), 231-238.

- Kelly, J.F., & Moos, R. (2003). Dropout from 12-step self-help groups: prevalence, predictors, and counteracting treatment influences. *Journal of Substance Abuse Treatment*, 24(3), 241-50.
- Kelly, J.F., Myers, M.G., & Brown, S.A. 2001. Do adolescents affiliate with 12-step groups? A multivariate process model of effects. *Journal of Studies on Alcohol*, 63, 293-304.
- Montgomery, H. A., Miller, W. R., & Tonigan, J. S. (1995). Does Alcoholics Anonymous involvement predict treatment outcome? *Journal of Substance Abuse Treatment*, 12(4), 241-246.
- White, W. & Hagen, R. (2005). Treatment, recovery, community: A call for reconnection. *Counselor*, 6(6), 52-56.