

The Status and Future of Addiction Recovery Support Services in the United States

William White, MA^a ; Keith Humphreys, PhD^b ; Monique Bourgeois, M.A^c ; Page Chiapella, PhD^d ; Arthur Evans, PhD^e ; Michael Flaherty, PhD^f ; Peter Gaumont, PhD^g ; Patrick Haggerson, M.A. ^h ; Beverley Haberle, M.H.Sⁱ ; Tom Hill, M.S.W.^j ; Lee Ann Kaskutas, PhD^k ; John Kelly, PhD^l ; Carol McDaid^m ; Joe Powell, L.D.C.ⁿ ; Christy Scott, Ph.D.^o ; Pat Taylor, M.S.W.^j

^a *Chestnut Health Systems, Bloomington, IL*

^b *Veterans Affairs and Stanford University Medical Centers, Palo Alto, CA*

^c *Association of Recovery Schools, Richmond, MN*

^d *National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD*

^e *Philadelphia Department of Behavioral Health, Philadelphia, PA;*

^f *Institute for Research, Education and Training in the Addictions, Jamison, PA;*

^g *Office of National Drug Control Policy, Gaithersburg, MD;*

^h *Betty Ford Institute, Palm Desert, CA*

ⁱ *The Council of Southeast Pennsylvania, Jamison, PA*

^j *Faces & Voices of Recovery, Washington, DC*

^k *Alcohol Research Group (ARG), Emeryville, CA*

^l *Harvard Center for Addiction Medicine, Boston, MA*

^m *Capitol Decisions, Washington, DC*

ⁿ *Associations of Persons Affected by Addiction, Dallas, TX*

^o *Chestnut Health Systems, Chicago, IL*

Authors Note: Any opinions expressed in this paper do not necessarily reflect the official policy positions of government agencies that employ or are advised by the authors. Address correspondence to: Keith Humphreys, PhD, Department of Psychiatry, 401 N. Quarry Road (MC: 5717), Stanford University School of Medicine, Stanford, CA 94305-5717. FAX (650) 617-2736.

Abstract

Addiction recovery support services (RSS), including recovery housing, schools, coaches, ministries, and community centers, are rapidly spreading in the United States. In February 2012, the Betty Ford Institute and the University of California, Los Angeles, convened an expert consensus conference to address four questions related to the growing RSS phenomenon: (1) What is new in the delivery of RSS in the United States? (2) What is known about RSS from the standpoint of science, practice, and experience? (3) What does the addiction treatment and recovery field need to know about RSS? (4) What are the next steps needed to develop, refine, and evaluate RSS? This article summarizes the expert panel's findings and recommendations.

Keywords: peer support, recovery, mutual aid, recovery support services, consensus conference

Introduction

Recovery from addiction is a voluntarily maintained lifestyle characterized by sobriety, personal health, and citizenship (Betty Ford Institute Consensus Panel, 2007). Historically, specialized support for recovery from alcohol and other drug addictions in the United States has been provided by peer-led mutual-aid organizations and professionally led addiction treatment organizations. The former include Native American abstinence-based religious/cultural revitalization and healing movements, 19th century recovery societies (e.g., Washingtonians, recovery-focused fraternal temperance societies, ribbon reform clubs, Keeley Leagues), Alcoholics Anonymous (AA) and its 12-step offshoots (e.g., Narcotics Anonymous, Cocaine Anonymous), mutual help organizations that reject any AA-style spiritual content (e.g., SMART Recovery) and, conversely, others (e.g., Celebrate Recovery) that forgo AA's generic "higher power" concept to instead employ an explicitly religious approach (Kelly & White, 2012). The latter span early inebriate homes and asylums, private addiction cure institutes, bottled and boxed home cures, and the late 20th century evolution of today's addiction treatment system in which evidence-based practices play an increasing role (White, 1998).

In the past two decades, a type of addiction recovery support has emerged that does not fit perfectly in either the category of peer-led recovery organizations or professional addiction treatment. This service form encompasses new social settings (e.g., recovery community organizations, recovery community centers, recovery homes, recovery schools, recovery industries, recovery ministries) and service roles (variably called recovery coaches/guides/mentors, recovery support specialists, or peer support specialists; Cousins, Antonini & Rawson, 2012; White, 2008, 2009; White, Kelly, & Roth, 2012).

The development of such "recovery support services" (RSS) raises numerous questions, opportunities, and challenges for the field. Accordingly, in February of 2012, the Betty Ford Institute (BFI), in collaboration with the University of California, Los Angeles (UCLA), hosted a consensus conference to explore the history, status, and future of recovery support services in the United States. Discussions were organized around four broad questions:

- 1) What is new in the delivery of recovery support services (RSS) in the United States?
- 2) What is known about RSS from the standpoint of science, practice, and experience?
- 3) What does the addiction treatment/recovery field need to know about RSS?
- 4) What are the next steps needed to develop, refine, and evaluate RSS?

This article summarizes the conclusions drawn from the BFI/UCLA Consensus Conference and briefly discusses the potential historical significance of RSS.

Several important historical trends set the stage for the 2012 BFI/UCLA Consensus Conference. These include the growth and diversification of addiction recovery experiences across and within secular, spiritual, and religious addiction recovery mutual-aid groups in the United States during the second half of the 20th century (Humphreys, 2004; White & Kurtz, 2006), the perceived disconnection between addiction treatment and the larger and more enduring process of long-term addiction recovery (Elise, 1999; Morgan, 1994), the emergence of a new grassroots recovery advocacy movement in the United States (White, 2007b), and the rise and maturation of recovery community organizations (RCOs) organized by and on behalf of

people in recovery, whose missions include policy advocacy and the delivery of peer-based recovery support services (Valentine, 2011; White, 2009, 2010).

Also influencing the emergence of RSS are the reconceptualization of addiction as a chronic medical disorder (Dennis & Scott, 2007; Flaherty, 2006; McLellan, 2002; McLellan, Lewis, O'Brien, & Kleber, 2000), the clearer articulation of the stages of long-term personal and family recovery (Brown & Lewis, 1999; Dennis, Scott, Funk, & Foss, 2005), calls to extend addiction treatment from models of acute biopsychosocial stabilization to models of sustained recovery management (Flaherty, 2006; Humphreys & Tucker, 2002), efforts to nest recovery management approaches within larger recovery-oriented systems of care (ROSC; Kelly & White, 2011; White, 2007a, 2008), and the promotion of RSS within and entirely outside of the formal health care system (Achara-Abrahams, Evans, & King, 2011; Clark, 2008; Kaplan, 2008; Kirk, 2011; Valentine, 2011). A further recent contextual influence is the expectation that services for alcohol and other drug problems, including RSS, will be expanded and integrated into other community service institutions as an outcome of the Affordable Care Act of 2010 (Buck, 2011; Hill, McDaid, & Taylor, 2012).

Seen as a whole, these trends reflect an increasing momentum toward the adoption of a recovery paradigm as a central organizing construct for the addictions field and the integration of RSS as part of larger transformation efforts within recovery-focused systems of care in the United States and in the United Kingdom (Berridge, 2012; El-Guebaly, 2012; White, 2005, 2008). These trends also suggest the need for a recovery research agenda to guide this re-orientation of the field's conceptual foundation and service practices (Laudet, Flaherty, & Langer, 2009; Laudet & Humphreys, 2012).

The Consensus Process

The Betty Ford Institute (BFI)—the prevention, education, and research arm of the Betty Ford Center—has for the past 7 years hosted a series of consensus conferences aimed toward the goals of defining recovery (Betty Institute Consensus Panel, 2007, McLellan, 2010), extending the benefits of addiction treatment through innovative continuing care strategies, (McKay et al., 2009), enhancing graduate medical school education in addiction (O'Connor, Nyquist, & McLellan, 2011), and breaking the intergenerational cycle of addiction through parent-focused strategies.

A total of 30 individuals participated in the RSS Consensus Conference convened in February 2012 at the Betty Ford Center in Rancho Mirage, California. The purpose of the conference was to discuss the status and future of RSS within the addictions field. Those present included a moderator, four speakers, three policy experts, seven community content experts (including representatives from recovery community organizations), three addiction research experts, and 15 observers. The policy experts included representatives from the National Institute on Alcohol Abuse and Alcoholism, the National Institute on Drug Abuse, and the Office of National Drug Control Policy.

The consensus process involved the presentation of papers on the history of RSS (White, 2012), the current RSS landscape (Hill et al., 2012), and research to date on RSS (Laudet & Humphreys, 2012); a facilitated discussion of the four guiding questions outlined above; and identification of points of group consensus and recommendations. To avoid premature

consensus, the facilitator and participants put special emphasis on drawing out disagreements and difficult issues.

Following the conference, participants were asked to complete an online survey rating their top RSS priorities drawn from the recommendations generated at the conference. Results were tabulated and are presented in the present article.

Results

There was clear consensus on the most significant aspects of RSS as they are developing in the United States. First, RSS rest on the premise that efforts to facilitate recovery from addiction should be grounded in an understanding of the chronicity and complexity of such disorders. A given RSS might be delivered in tandem with treatment, mutual aid organizations, or both of these entities, or it might be delivered independently. As a class of intervention, however, RSS are organized to offer longer-term support than that provided by acute care addiction treatment as well as a broader range of support than that provided by either recovery mutual-aid organizations or addiction treatment programs.

Second, RSS are distinctive in their frequent delivery within new recovery-focused entities (recovery community organizations, recovery homes, recovery schools, recovery industries, recovery ministries, recovery cafes) and within such varied community settings as jails, schools, and Federally Qualified Health Centers. Rather than exclusively ask, “How do we get the person in need to a treatment center or mutual aid meeting?” RSS providers ask, “How do we deliver nested recovery support within the settings where he or she is already involved?” As this conceptualization suggests, the formal treatment system is only one of many venues through which RSS are provided.

Third, those providing RSS extol the legitimacy of multiple long-term pathways of recovery and value personal choice in the recovery process. This distinguishes them from mutual help organizations that facilitate travel down a single pathway to recovery (i.e., the pathway of that particular fellowship). RSS include assertive linkage to a broad spectrum of recovery mutual aid and treatment options, provide a larger menu of recovery support options, and utilize recovery-focused service planning protocols quite different from traditional treatment planning schema (Borkman, 1997).

Fourth, involvement of persons in recovery is common in the development and delivery of RSS (White, 2009). At the same time, individuals who have not personally experienced addiction are also often involved in RSS.

Fifth, RSS are distinctive in offering support across the stages of recovery—spanning pre-recovery engagement and recovery priming, recovery initiation and stabilization, recovery maintenance, enhanced quality of personal and family life in long-term recovery, and support in breaking intergenerational cycles of problem transmission. Also of note is that RSS can be provided before, after, or in tandem with professional treatment, or in lieu of such treatment. Prior or concurrent treatment is rarely a requirement for participation in RSS.

Other distinguishing aspects of RSS noted in the consensus panel discussions include the more assertive style of RSS (compared to most recovery mutual aid or professional treatment initiatives), the greater emphasis on advocacy as a critical dimension of RSS, the extensive use of volunteers—particularly peers—in the delivery of RSS, the differences in ethical guidelines

for RSS compared to those for addiction counselors, and a focus on the community as an object of change as well as the individual and family (White, 2012).

In reviewing existing RSS trends and RSS-related research, the consensus panel drew five major conclusions.

- 1) RSS have been implemented in a wide variety of organizational and community settings (Hill et al., 2012). Further, RSS have been tailored to the needs of special populations (including support for those in medication-assisted recovery from opioid addiction), adapted across diverse cultural and linguistic contexts, and often provided in gender-based settings.
- 2) Experience confirms that there is demand for RSS. Individuals and families in all stages of recovery are seeking and using such services in tandem with, and in lieu of, professional treatment and mutual help organizations.
- 3) Recovery community organizations (RCOs) and other organizations are identifying critical aspects of the implementation of RSS, e.g., methods of recruiting, screening, training, and supervising RSS providers.
- 4) The proliferation of RSS in the United States and the United Kingdom is far ahead of the research directly evaluating the short- and long-term effects of such services (Laudet & Humphreys, 2012.) Nevertheless, the theoretical rationale for RSS is buttressed by research showing the positive influence of social support on short- and long-term recovery outcomes (Beattie & Longabaugh, 1999; Groh, Jason, Davis, Olson, & Ferrari, 2007; Humphreys, Mankowski, Moos, & Finney, 1999; Humphreys, Moos, & Cohen, 1997; Stout, Kelly, Magill & Pagano, 2012), comparisons of social versus medical model programs (Kaskutas, Witbrodt, & French, 2004; Kaskutas, Zavala, Parthasarathy, & Witbrodt, 2008), and studies evaluating assertive approaches to recovery checkups and continuing care for adults (Dennis & Scott, 2012; McKay, 2005, 2009; Scott & Dennis, 2009; Scott, Dennis, & Foss, 2005) and adolescents (Godley, Godley, Dennis, Funk, & Passetti, 2007).
- 5) Research in some RSS arenas is quite strong (e.g., recovery housing; Jason & Ferrari, 2010; Jason et al., 2007; Polcin, Korcha, Bond, & Galloway, 2010). For others, the research base is descriptive (e.g., recovery schools and collegiate recovery communities; Cleveland, Harris, Baker, Herbert, & Dean, 2007; Harris, Baker, Kimball, & Shumway, 2008; Moberg & Finch, 2008) or non-existent (e.g., recovery community centers, recovery-focused work programs, recovery ministries, and recovery cafés). Especially troubling is the paucity of research on RSS specialty roles (e.g., recovery coaches), given their rapid proliferation across the United States as a component of efforts to embrace models of recovery management and recovery-oriented systems of care. More positively, administrative data gathered on high-profile national initiatives, such as the Substance Abuse and Mental Health Services Administration's Recovery Community Services Program and Access to Recovery, suggest positive effects of RSS on recovery outcomes (e.g., sobriety/remission, improvements in global health, community reintegration; Laudet & Humphreys, 2012; White, 2009).

Consensus conference discussions of an RSS research agenda focused, in order of perceived priority, on the need to (1) assess the effectiveness and cost-effectiveness of particular types of RSS in producing outcomes in individuals and families, within treatment systems,

within local communities of recovery, and within communities more broadly; (2) gather and synthesize practice-based evidence and wisdom related to the optimal design and delivery of RSS; (3) conduct surveys that quantify the magnitude and methods being used to deliver RSS in the United States; (4) identify active ingredients of RSS and particular service combinations and sequences that most strongly influence recovery outcomes; and (5) create and validate instruments to assess individual and family needs for particular types of RSS. Some of the more specific questions posed within this discussion are presented in Table 1.

Table 1: What we need to know about Recovery Support Services (RSS)

<p>Assessing the effectiveness and cost-effectiveness of RSS</p>	<p>What are key elements of recovery that would lend themselves to measurement as RSS outcomes (e.g., sense of self-efficacy, quality of life, recovery/social capital, etc.)?</p> <p>Are there differences in outcomes influenced by the recovery status of the person delivering RSS or by the organizational setting through which RSS are delivered?</p> <p>Does the duration and amount of contact influence the effectiveness of RSS?</p> <p>Do RSS avert or reduce use of costly acute care services (e.g., emergency rooms)?</p> <p>What existing measures might be adopted to measure some or all identified elements?</p>
<p>Optimal design for RSS</p>	<p>What is a typical menu of RSS?</p> <p>How do RSS coming out of RCOs and treatment organizations differ from each other, if at all, and from those being offered by the life coaching field?</p> <p>What are the key quality indicators of RSS?</p> <p>What qualifications are required to deliver RSS? How does personal or family recovery by providers affect delivery of RSS?</p> <p>Does the impact of RSS vary depending on whether staff are volunteers, paid, or a combination of such, or in recovery, not in recovery, or a combination?</p> <p>How are RSS being refined across boundaries of age, gender, culture, drug choice, and pathways of recovery?</p>

	<p>Is there an optimal length or intensity of service that RSS should provide?</p> <p>How many people can a single RSS provider serve simultaneously in different contexts and with different populations?</p> <p>How can people providing RSS be best supervised?</p> <p>How can RSS avoid becoming either a cheap form of treatment and a rationale for cutting treatment services, or, becoming commercialized to the point that they lose their grassroots nature and commitment to advocacy that make them appealing to those who access them ?</p> <p>What are the critical considerations in developing new RSS and new RSS service locations?</p>
<p>National, state, and local assessment of currently available RSS</p>	<p>What is the present national, state, and local availability of RSS?</p> <p>What is the current need/demand for particular types of RSS?</p> <p>What is the RSS needs profile of people at different stages of recovery?</p> <p>How is technology being utilized in the delivery of RSS?</p>
<p>Active ingredients/combinations of RSS</p>	<p>What are the most potent ingredients of RSS in terms of their influence on recovery outcomes?</p> <p>Which individuals are most likely to benefit from RSS?</p> <p>Are there particularly potent RSS combinations or sequences? Do these combinations or sequences need to be implemented during certain “critical” recovery phases to be effective?</p> <p>Are there ingredients unique to RSS that are not found in recovery mutual aid or professional addiction treatment?</p>
<p>Assessing individual and family need for RSS</p>	<p>Who is most likely to need and utilize RSS?</p> <p>How are organizations delivering RSS evaluating the support needs and service response of those they serve, e.g., individual and community needs assessments, tools, checklists, recovery planning formats?</p> <p>Is there a needs-and-strengths profile of people at different stages of</p>

	recovery? If not, could it be developed?
--	--

The closing discussion of the RSS Consensus Conference generated an action agenda for RSS service providers, policymakers and administrators, and the research community. Table 2 displays the specific recommendations within these three arenas.

Table 2: Recommended and Prioritized Next Steps in the Evolution of RSS

<p>For the RSS practice community</p>	<p>Clarify role boundaries between RSS provider, addiction counselor, life coach, and other service roles (e.g., case managers, outreach workers) and the relationship between RSS, recovery mutual aid organizations, and professional addiction treatment.</p> <p>Communicate to multiple stakeholders the precise purpose of RSS and their role in the larger service/support system and in the broader community.</p> <p>Survey RCOs, treatment providers, Access to Recovery sites, and Single State Agencies to establish a benchmark of current RSS practices.</p> <p>Create a central repository of documents, tools, media, etc., that can be used to enhance replication of RSS.</p> <p>Disseminate documents that can help educate policymakers, funders, treatment and allied service professionals, and the recovery community about the nature of RSS and evaluation results.</p>
<p>Policymakers and administrators</p>	<p>Integrate RSS within the implementation of the Affordable Care Act.</p> <p>Integrate questions related to RSS in national surveys of recovery prevalence and addiction treatment.</p> <p>Provide technical assistance and support to enhance the mobilization of local recovery communities and development of local recovering community organization RSS capabilities.</p> <p>Host and participate in conferences that include an overview of federal RSS initiatives, local RCO activities related to RSS, and what private managed behavioral health companies and disease management programs are doing with RSS.</p> <p>Evaluate states' Medicaid experiences with RSS and disseminate the</p>

	findings
Researchers	<p>Pursue quantitative and qualitative outcome studies of RSS.</p> <p>Develop a valid and reliable measure of recovery</p> <p>Use existing data sets to assess the potential need/market for RSS.</p> <p>Design new research in light of findings of prior studies (e.g. Kaskutas et al., 2004).</p> <p>Develop standard evaluation protocols for use by organizations delivering RSS.</p> <p>Create opportunities for dialogue between addiction researchers and representatives from recovery community organizations involved in RSS delivery.</p> <p>Examine comparable RSS for other health conditions for their effectiveness and transferability to addiction RSS</p>

Discussion

The BFI/UCLA Consensus Conference review of current research and prevailing practices related to recovery support services drew five broad conclusions.

1. RSS are being implemented in diverse geographical, cultural, and organizational settings for people seeking recovery from addiction, those newly in recovery, and those in long-term recovery. Some people access RSS in combination with addiction treatment programs and mutual help organizations, whereas others do so entirely independent of these services.
2. Payers, providers, and the recovery community are showing increasing interest in RSS as a way to support recovery from addiction.
3. Although the development of RSS has strong theoretical support, rigorous research has been limited to particular recovery support institutions (e.g., recovery residences) and to particular recovery support protocols (e.g., posttreatment recovery management checkups).
4. The expansion of existing RSS, replication of RSS in new settings, and development of new RSS need to be informed by studies evaluating the effects of a wide variety of RSS components on short- and long-term recovery outcomes for individuals and families, and studies that contribute to the development of RSS practice guidelines and standards. At the same time, the practice of RSS cannot wait to proceed until traditional academic research projects are completed. Researchers and providers should gather, synthesize, and attend to practice-based evidence, rather than assuming that traditional research approaches can be the only guide to action.

5. The further development, evaluation, and refinement of RSS in the United States will require concerted action by the RSS practice community, policymakers, systems administrators, and the research community.

There are a number of important ways in which RSS models being replicated in U.S. communities could expand support for individuals recovering from addiction. RSS providers, such as recovery mutual-aid groups, frequently draw on the rich experiential knowledge and expertise of individuals in recovery (Borkman, 1976; Jackson, 2001). RSS could provide a menu of support services to a broad spectrum of individuals (including those without past or current treatment involvement) for an extended period of time (including pre-recovery engagement and long-term personal or family recovery). RSS can expand access to services and supports because they are delivered primarily within the natural environments of those receiving them (homes, neighborhoods, schools, churches, hospitals, jails). Another distinctive feature about RSS is that they strive for change not just in individuals but in the community. Specifically, RSS attempt to develop community recovery capital—the physical, psychological, and social space within the community where recovery can flourish (Granfield & Cloud, 1999; Humphreys et al., 1997). This focus on community goes well beyond the traditional clinical perspective of changing individuals (or, at most, individuals and their immediate family) and reflects the advocacy component of RSS.

RSS serve individuals who are on diverse pathways of recovery, including persons who initiate and sustain recovery without involvement in a mutual aid group. The RSS provider must therefore become knowledgeable and fluent across multiple pathways and styles of recovery initiation and maintenance, including working with persons with lower problem severity/complexity. RSS also address issues beyond initiation and maintenance of recovery, e.g., housing, education, employment, debt management, criminal record expungement, physical and mental health, exercise and nutrition, social networking, community service, advocacy opportunities, and recovery-linked involvement in art, literature, music, theatre, or sports.

RSS represent a new menu of recovery support mechanisms delivered in natural community environments through novel social institutions (e.g., recovery community organizations [RCOs]) and innovative service roles (e.g., recovery coach). The RCO is distinct from treatment organizations and recovery mutual-aid groups, and the recovery coach fits neither the categories of addiction counselor nor recovery mutual-aid sponsor/mentor. The critical question regarding the ideal means of resolving alcohol and other drug problems is not, “Should people seeking recovery choose professional treatment or a recovery mutual-aid group or RSS?” It is rather, “Are there ways that professional treatment, recovery mutual-aid societies, and RSS can be created and organized such that more numerous and more diverse long-term recovery pathways are available to more people?” Natural community experiments and controlled trials are needed to discover, for the treatment of addiction, what combined, sequenced, and stand-alone interventions have discovered for the treatment of AIDS and cancer. The advent of RSS creates an historic opportunity for such discoveries, though considerable challenges lie ahead in more clearly defining and more rigorously evaluating this expanding menu of services delivered through new roles and in new service settings.

Acknowledgment: The 2012 BFI/UCLA Consensus Conference on Recovery Support Services was financially supported by Norlien Foundation, Hilton Foundation, and American Legacy

Foundation (Legacy for Health). We thank Richard A. Rawson and Wilson Compton for very helpful critiques of these ideas. The RSS Consensus Conference was facilitated by Keith Humphreys, with Sarah J. Cousins and Kristi Wellenberger serving as recorders.

References

- Achara-Abrahams, I., Evans, A. C., & King, J. K. (2011). Recovery-focused behavioral health systems transformation: A framework for change and lessons learned from Philadelphia. In J. F. Kelly & W. L. White (Eds.), *Addiction recovery management: Theory, science and practice* (pp. 187–208). New York: Springer Science.
- Beattie, M. C., & Longabaugh, R. (1999). General and alcohol-specific social support following treatment. *Addictive Behaviors, 24*, 593–606.
- Berridge, V. (2012). The art of medicine: The rise, fall, and revival of recovery in drug policy. *Lancet, 379*, 22–23.
- Betty Ford Institute Consensus Panel. (2007). What is recovery? A working definition from the Betty Ford Institute. *Journal of Substance Abuse Treatment, 33*, 221–228.
- Borkman, T. (1976). Experiential knowledge: A new concept for the analysis of self-help groups. *Social Service Review, 50*, 443–456.
- Borkman, T. J. (1997). Is recovery planning any different from treatment planning? *Journal of Substance Abuse Treatment, 15*(1), 37–42.
- Brown, S., & Lewis, B. (1999). *The alcoholic family in recovery: A developmental model*. New York: Guilford.
- Buck, J. A. (2011). The looming expansion and transformation of public substance abuse treatment under the Affordable Care Act. *Health Affairs, 30*(8), 1402–1410.
- Clark, W. (2008). *Recovery-oriented systems of care: SAMHSA/CSAT's public health approach to substance use problems & disorders. Aligning concepts, practice, and contexts to promote long term recovery: An action plan*. Philadelphia, PA: Institute for Research, Education & Training in Addictions (IRETA).
- Cleveland, H. H., Harris, K. S., Baker, A. K., Herbert, R., & Dean, L. R. (2007). Characteristics of a collegiate recovery community: Maintaining recovery in an abstinence-hostile environment. *Journal of Substance Abuse Treatment, 33*(1), 13–23.
- Cousins, S. J., Antonini, V. P., & Rawson, R. A. (2012). Utilization, measurement, and funding of recovery support services. *Journal of Psychoactive Drugs, 44* (4), 325–333.
- Dennis, M. L., & Scott, C. K. (2007). Managing addiction as a chronic condition. *Addiction Science & Clinical Practice, 4*(1), 45–55.
- Dennis, M. L., & Scott, C. K. (2012). Four-year outcomes from the Early Re-Intervention (ERI) experiment using Recovery Management Checkups (RMCs). *Drug and Alcohol Dependence, 121*(1–2), 10–7.
- Dennis, M. L., Scott, C. K., Funk, R., & Foss, M. A. (2005). The duration and correlates of addiction and treatment careers. *Journal of Substance Abuse Treatment, 28*, S51–S62.
- El-Guebaly, N. (2012). The meanings of recovery from addiction: Evolution and promises. *Journal of Addiction Medicine, 6*(1), 1–9.
- Elise, D. (1999). Recovering recovery. *Journal of Ministry in Addiction and Recovery, 6*(2), 11–23.

- Flaherty, M. (2006). *Special Report: A unified vision for the prevention and management of substance use disorders: Building resiliency, wellness and recovery – A shift from an acute care to a sustained care recovery management model*. Pittsburgh, PA.: Institute for Research, Education & Training in Addictions (IRETA).
- Godley, M. D., Godley, S. H., Dennis, M. L., Funk, R. R., & Passetti, L. L. (2007). The effect of assertive continuing care on continuing care linkage, adherence, and abstinence following residential treatment for adolescent substance use disorders. *Addiction, 102*, 81–93.
- Granfield, R., & Cloud, W. (1999). *Coming clean: Overcoming addiction without treatment*. New York: New York University Press.
- Groh, D. R., Jason, L. A., Davis, M. I., Olson, B. D., & Ferrari, J. R. (2007). Friends, family, and alcohol abuse: An examination of general and alcohol-specific social support. *The American Journal on Addictions, 16*, 49–55.
- Harris, K., Baker, A., Kimball, T., & Shumway, S. (2008). Achieving systems-based sustained recovery: A comprehensive model for collegiate recovery communities. *Journal of Groups in Addiction and Recovery, 2*(2–4), 220–237.
- Hill, T., McDaid, C., & Taylor, P. (2012, February). *Peer recovery support services: Evolving community-based practices and infrastructure*. Presented at the 2012 Betty Ford Institute/UCLA annual recovery conference, February 21–23, Rancho Mirage, CA.
- Humphreys, K. (2004). *Circles of recovery: Self-help organizations for addictions*. Cambridge: Cambridge University Press.
- Humphreys, K., Mankowski, E., Moos, R. H., & Finney, J. W. (1999). Do enhanced friendship networks and active coping mediate the effect of self-help groups on substance use? *Annals of Behavioral Medicine, 21*(1), 54–60.
- Humphreys, K., Moos, R. J., & Cohen, C. (1997). Social and community resources and long-term recovery from treated and untreated alcoholism. *Journal of Studies on Alcohol, 58*(3), 231–238.
- Humphreys, K., & Tucker, J. (2002). Towards more responsive and effective intervention systems for alcohol-related problems. *Addiction, 97*, 126–132.
- Jackson, S. W. (2001). The wounded healer. *Bulletin of the History of Medicine, 75*, 1–36.
- Jason, L. A., & Ferrari, J. R. (2010). Oxford House recovery homes: Characteristics and effectiveness. *Psychological Services, 7*(2), 92–102.
- Jason, L. A., Olson, B. D., Ferrari, J. R., Majer, J. M., Alvarez, J., & Stout, J. (2007). An examination of main and interactive effects of substance abuse recovery housing on multiple indicators of adjustment. *Addiction, 102*(7), 1114–1121.
- Kaplan, L. (2008). *The role of recovery support services in recovery-oriented systems of care* (DHHS Publication No. (SMA) 08-4315). Rockville, MD: Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration.
- Kaskutas, L. A., Witbrodt, J., & French, M. T. (2004). Outcomes and costs of day hospital treatment and nonmedical day treatment for chemical dependency. *Journal of Studies of Alcohol, 65*(3), 371–382.
- Kaskutas, L. A., Zavala, S. K., Parthasarathy, S., & Witbrodt, J. (2008). Costs of day hospital and community residential chemical dependency treatment. *Journal of Mental Health Policy Economics, 11*(1), 27–32.
- Kelly, J. F., & White, W. L. (Eds.) (2011). *Addiction recovery management: Theory, science and practice*. New York: Springer Science.

- Kelly, J. F., & White, W. L. (2012). Broadening the base of addiction mutual help. *Journal of Groups in Addiction Recovery*, 7, (2–4), 82–101.
- Kirk, T. (2011). Connecticut's journey to a statewide recovery-oriented health-care system: Strategies, successes, and challenges. In J. F. Kelly, & W. L. White (Eds.), *Addiction recovery management: Theory, science and practice* (pp. 209–235). New York: Springer Humana Press.
- Laudet, A., Flaherty, M., & Langer, D. (2009). *Building the science of recovery*. Pittsburgh, PA: Institute for Research, Education and Training and Northeast Addiction Technology Transfer Center.
- Laudet, A. B., & Humphreys, K. (2012, February). *Promoting recovery in an evolving context: What we know and what we need to know about recovery support services*. Presented at the 2012 BFI/UCLA annual recovery conference, February 21–23, Rancho Mirage, CA.
- McKay, J. R. (2005). Is there a case for extended interventions for alcohol and drug use disorders? *Addiction*, 100(11), 1594–1610.
- McKay, J. R. (2009). Continuing care research: What we have learned and where we are going. *Journal of Substance Abuse Treatment*, 36(2), 131–145.
- McKay, J. R., Carise, D., Dennis, M. L., Dupont, R., Humphreys, K., Kemp, J.,...Schwartzlose, J. (2009). Extending the benefits of addiction treatment: Practical strategies for continuing care and recovery. *Journal of Substance Abuse Treatment*, 36, 127–130.
- McLellan, A. T. (2002). Have we evaluated addiction treatment correctly? Implications from a chronic care perspective. *Addiction*, 97(3), 249–252.
- McLellan, A. T. (2010). What is recovery? Revisiting the Betty Ford Institute Consensus Panel definition: The Betty Ford Consensus Panel and consultants. *Journal of Substance Abuse Treatment*, 38, 200–201.
- McLellan, A. T., Lewis, D. C., O'Brien, C. P., & Kleber, H. D. (2000). Drug dependence, a chronic medical illness: Implications for treatment, insurance, and outcomes evaluation. *Journal of the American Medical Association*, 284(13), 1689–1695.
- Moberg, D. P., & Finch, A. J. (2008). Recovery high schools: A descriptive study of school programs and students. *Journal of Groups in Addiction and Recovery*, 2, 128–161.
- Morgan, O. J. (1994). Extended length of sobriety: The missing variable. *Alcoholism Treatment Quarterly*, 12(1), 59–71.
- O'Connor, P. G., Nyquist, J. G., & McLellan, A. T. (2011). Integrating addiction medicine into graduate medical education in primary care: The time has come. *Annals of Internal Medicine*, 154(1), 56–60.
- Polcin, D. L., Korcha, R. A., Bond, J., & Galloway, G. (2010). Sober living houses for alcohol and drug dependence: 18-month outcomes. *Journal of Substance Abuse Treatment*, 38(4), 356–365.
- Scott, C. K., & Dennis, M. L. (2009). Results from two randomized clinical trials evaluating the impact of quarterly recovery management checkups with adult chronic substance users. *Addiction*, 104(6), 959–971.
- Scott, C. K., Dennis, M. L., & Foss, M. A. (2005). Utilizing recovery management checkups to shorten the cycle of relapse, treatment re-entry, and recovery. *Drug and Alcohol Dependence*, 78, 325–338.
- Stout, R. L., Kelly, J. F., Magill, M., & Pagano, M. E. (2012) Association between social influences and drinking outcomes across three years. *Journal of Studies on Alcohol and*

- Drugs*, 73(3), 489–497.
- Valentine, P. (2011). Peer-based recovery support services within a recovery community organization: The CCAR experience. In J. F. Kelly & W. L. White (Eds.), *Addiction recovery management: Theory, science and practice* (pp. 259–279). New York: Springer Science.
- White, W. L. (1998). *Slaying the dragon: The history of addiction treatment and recovery in America*. Bloomington, IL: Chestnut Health Systems.
- White, W. L. (2005). Recovery: Its history and renaissance as an organizing construct concerning alcohol and other drug problems. *Alcoholism Treatment Quarterly*, 23(1), 3–15.
- White, W. L. (2007a). *Perspectives on systems transformation: How visionary leaders are shifting addiction treatment toward a recovery-oriented system of care. (Interviews with H. Westley Clark, Thomas A. Kirk, Jr., Arthur C. Evans, Michael Boyle, Phillip Valentine and Lonnetta Albright)*. Chicago, IL: Great Lakes Addiction Technology Transfer Center.
- White, W. L. (2007b). The new recovery advocacy movement in America. *Addiction*, 102, 696–703.
- White, W. L. (2008). Recovery: Old wine, flavor of the month or new organizing paradigm? *Substance Use and Misuse*, 43(12–13), 1987–2000.
- White, W. L. (2009). *Peer-based addiction recovery support: History, theory, practice, and scientific evaluation*. Chicago, IL: Great Lakes Addiction Technology Transfer Center and Philadelphia Department of Behavioral Health and Mental Retardation Services.
- White, W. L. (2010). Non-clinical addiction recovery support services: History, rationale, models, potentials and pitfalls. *Alcoholism Treatment Quarterly*, 28, 256–272.
- White, W. L. (2012, February). *Historical perspectives on addiction recovery support*. Presented at the 2012 BFI/UCLA annual recovery conference, February 21–23, Rancho Mirage, CA.
- White, W., Kelly, J., & Roth, J. (2012). New addiction recovery support institutions: Mobilizing support beyond professional addiction treatment and recovery mutual aid. *Journal of Groups in Addiction & Recovery*, 7(2–4), 297–317.
- White, W., & Kurtz, E. (2006). The varieties of recovery experience. *International Journal of Self Help and Self Care*, 3(1–2), 21–61.